


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C07D 405/12, A61K 31/445	A1	(11) International Publication Number: WO 96/16060 (43) International Publication Date: 30 May 1996 (30.05.96)
(21) International Application Number: PCT/EP95/04519 (22) International Filing Date: 16 November 1995 (16.11.95) (30) Priority Data: 94203421.6 24 November 1994 (24.11.94) EP (34) Countries for which the regional or international application was filed: DE et al. (71) Applicant (for all designated States except US): JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B-2340 Beerse (BE). (71) Applicants (for US only): HENDRICKX, Marie-Louise (heirress of the deceased inventor) [BE/BE]; Kongostraat 156, B-2300 Turnhout (BE). VAN DAELE, Kurt, Godfried, Cornelius, Emile (heir of the deceased inventor) [BE/BE]; Berchemlei 213, B-2140 Borgerhout (BE). VAN DAELE, Peter, Jules, Victor (heir of the deceased inventor) [BE/BE]; Berkenlaan 54A, B-1850 Grimbergen (BE). VAN DAELE, Glenn, Kurt, Ludo (heir of the deceased inventor) [BE/BE]; Steenweg op Oosthoven 110, bus 8, B-2300 Turnhout (BE). (72) Inventor: VAN DAELE, Georges, Henri, Paul (deceased).		(72) Inventors; and (75) Inventors/Applicants (for US only): BOSMANS, Jean-Paul, René, Marie, André [BE/BE]; Hofgracht 13, B-2650 Edegem (BE). SCHUURKES, Joannes, Adrianus, Jacobus [NL/BE]; Veldstraat 70, B-2340 Beerse (BE). (81) Designated States: AL, AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, LS, MW, SD, SZ, UG). Published <i>With international search report.</i>

(54) Title: ENTEROKINETIC BENZAMIDE**(57) Abstract**

The present invention is concerned with a novel benzamide derivative of formula (I). Pharmaceutical compositions comprising said novel compounds processes for preparing compounds and compositions, and the use thereof as a medicine, in particular in the treatment of conditions involving a decreased motility of the intestine are described.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LJ	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

ENTEROKINETIC BENZAMIDE

- The present invention is concerned with a novel benzamide derivative and the pharmaceutically acceptable acid addition salts thereof, pharmaceutical compositions comprising said novel compound, processes for preparing said compounds and compositions, and the use thereof as a medicine, in particular in the treatment of conditions involving an impaired motility of the intestine, especially of the colon.
- 5 In our EP-0,389,037-A, published on September 26, 1990, N-(3-hydroxy-4-piperidin-yl) (dihydrobenzofuran or dihydro-2H-benzopyran)carboxamide derivatives are disclosed as having gastrointestinal motility stimulating properties. In our EP-0,445,862-A, published on September 11, 1991, N-(4-piperidinyl) (dihydrobenzofuran or dihydro-2H-benzopyran)carboxamide derivatives are disclosed also having
- 10 gastrointestinal motility stimulating properties.
- 15

The compound subject to the present application differs therefrom by showing superior enterokinetic properties.

- 20 The present invention concerns a compound of formula



and the pharmaceutically acceptable acid addition salts thereof.

- 25 The chemical name of the compound of formula (I) is 4-amino-5-chloro-2,3-dihydro-N-[1-(3-methoxypropyl)-4-piperidinyl]-7-benzofurancarboxamide.

- The pharmaceutically acceptable acid addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic acid addition salt forms which the compounds of formula (I) are able to form. The latter can conveniently be obtained by
- 30 treating the base form with such appropriate acid. Appropriate acids comprise, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid; sulfuric; nitric; phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic, malonic, succinic (i.e. butane-

-2-

- dioic acid), maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pamoic and the like acids. The term addition salt as used hereinabove also comprises the solvates which the compounds of formula (I) as well as the salts thereof are able to form. Such solvates are, for example, hydrates, alcoholates and the like. Conversely the salt form can be converted by treatment with alkali into the free base form. Hereinafter the term "compounds of formula (I)" means the compound of formula (I) as well as the pharmaceutically acceptable acid addition salts thereof, unless otherwise mentioned.
- 10 Interesting compounds of formula (I) are the acid addition salts which are formed by treating the base form of the compound of formula (I) with hydrohalic acids or butanedioic acid.

- Preferred compounds of formula (I) are 4-amino-5-chloro-2,3-dihydro-N-[1-(3-methoxypropyl)-4-piperidinyl]-7-benzofurancarboxamide monohydrochloride and
- 15 4-amino-5-chloro-2,3-dihydro-N-[1-(3-methoxypropyl)-4-piperidinyl]-7-benzofuran-carboxamide butanedioate (1:1).


- The compounds of formula (I) may be prepared according to procedures which are disclosed in EP-0,389,037-A and EP-0,445,862-A. A number of preparation alternatives are shown hereinunder.
- 20

- In the following preparations, the reaction products may be isolated from the reaction mixture and, if necessary, further purified according to methodologies generally known in the art such as, for example, extraction, distillation, crystallization, trituration and chromatography.
- 25


- The compounds of formula (I) can be prepared by N-alkylating an intermediate of formula (II) with an alkylating reagent of formula (III), wherein W is an appropriate leaving group such as a halo, e.g. chloro; or a sulfonyloxy leaving group, e.g. methanesulfonyloxy (mesylate) or a *p*-toluenesulfonyloxy (tosylate) in a reaction inert solvent such as a dipolar aprotic solvent, e.g. dimethyl formamide, in the presence of an appropriate base such as, for instance, triethylamine. A suitable catalyst such as potassium iodide may also be added to enhance the reaction rate.
- 30

35

-3-



The compound of formula (I) may also be prepared by an N-acylation reaction of a carboxylic acid of formula (IV) or a reactive intermediate thereof and an amine of formula (V). Said N-acylation reaction may be performed by stirring the two reactants in a reaction-inert solvent, such as a chlorinated hydrocarbon, e.g. chloroform, or an aromatic hydrocarbon, e.g. toluene.



The above-mentioned intermediates are art-known or the preparation thereof is mentioned in EP-0,389,037-A and EP-0,445,862-A.

The compounds of formula (I) possess excellent intestinal motility stimulating properties. In particular the present compounds of formula (I) show significant motility enhancing effects on the small and large intestine. In other words the present compounds of formula (I) have enterokinetic properties. These properties are supported by the pharmacological examples described hereinunder. The present compounds of formula (I) enhance non-adrenergic non-cholinergic (NANC) excitation and the propulsion of faecal pellets through the large bowel. In addition, they accelerate gastric emptying and small intestinal contractile activity and have a facilitating effect on the cholinergic nerves. Said compounds are also devoid of 5-HT₂ or 5-HT₃ receptor antagonistic properties. Moreover, the present compounds also show *in-vivo* activity as is evidenced in the "Telemetric recording of colonic motility in conscious dogs" test.

In view of their useful enterokinetic enhancing properties the subject compounds may be formulated into various forms for administration purposes.

To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, in base or acid addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirably in unitary dosage form suitable, preferably, for administration orally, rectally or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not cause a significant deleterious effect to the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on, as an ointment.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used in the specification and claims herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such dosage unit forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, injectable solutions or suspensions, teaspoonfuls, tablespoonfuls and the like, and segregated multiples thereof.

In view of their capability to stimulate the motility of the intestinal system and in

particular their capacity to enhance the motility of the colon, the subject compounds are useful to normalize or to improve the intestinal transit in subjects suffering from symptoms related to disturbed motility, e.g. a decreased peristalsis of the small and large intestine alone or in combination with delayed gastric emptying.

- 5 In view of the utility of the compounds of the present invention, there is provided a method of treating warm-blooded animals, including humans, suffering from motility disorders of the intestinal system, such as, for example, constipation, pseudo-obstruction, intestinal atony, post-operative intestinal atony, irritable bowel syndrome (IBS), and drug-induced delayed transit. In particular there is provided a method of
- 10 treating large bowel motility disorders. The subject compounds may also be used to facilitate large bowel cleaning or to facilitate intubation and/or endoscopy. Said method comprises the systemic administration of an effective (small and large) intestinal stimulating amount of a compound of formula (I), to warm-blooded animals, including humans. Hence, the use of a compound of formula (I) as medicine is provided, and in
- 15 particular the use of a compound of formula (I) for the manufacture of a medicine for treating conditions involving a disordered motility or transit of the small and large intestine.

- In general it is contemplated that a therapeutically effective amount would be from about
- 20 0.001 mg/kg to about 10 mg/kg body weight, preferably from about 0.02 mg/kg to about 5 mg/kg body weight. A method of treatment may also include administering the active ingredient on a regimen of between two or four intakes per day.

25 Experimental part

Example 1

- In trichloromethane (135 ml) 4-amino-5-chloro-2,3-dihydro-7-benzofurancarboxylic acid (0.05 mol) (the preparation of which was described in EP-0,389,037-A) was suspended and cooled to $\pm 5^{\circ}\text{C}$. *N,N*-diethylethanamine (0.05 mol) was added dropwise at a
- 30 temperature below 10°C . Ethyl chloroformate (0.05 mol) was added dropwise and the reaction mixture was stirred for 40 min. while keeping the temperature below 10°C . The resulting mixture was added dropwise over a 20-min period to a solution of 1-(3-methoxypropyl)-4-piperidinamine (0.05 mol) in trichloromethane (35 ml). The cooling bath was removed and the reaction mixture was stirred for 150 min. Said
- 35 mixture was washed with water (50 ml). The precipitate was filtered off over a glass filter and washed with water and CHCl_3 . The filtrate was separated in its layers. The separated organic layer was washed with water (50 ml) + a 50% NaOH solution (1 ml),

- dried, filtered and the solvent was evaporated. The residue was stirred in 2-propanol (100 ml). This mixture was acidified with HCl/2-propanol (7.2 ml; 5.29 N). The mixture was stirred for 16 hours at room temperature and the resulting precipitate was filtered off, washed with 2-propanol (15 ml) and dried (vacuum; 50 °C), yielding 12.6 g (62%) of 4-amino-5-chloro-2,3-dihydro-N-[1-(3-methoxypropyl)-4-piperidinyl]-7-benzofurancarboxamide monohydrochloride (comp. 1).

Example 2

- A mixture of 4-amino-5-chloro-2,3-dihydro-N-(4-piperidinyl)-7-benzofurancarboxamide (0.01 mol), 1-chloro-3-methoxypropane (0.012 mol), N,N-diethylethanamine (2.1 ml) and KI (catalytic amount) in N,N-dimethylformamide (75 ml) was stirred overnight at 50 °C. The reaction mixture was cooled. The solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CHCl₃/(CH₃OH/NH₃) 97/3). The pure fractions were collected and the solvent was evaporated. The residue was dissolved in 2-propanol and converted into the hydrochloric acid salt (1:1) with HCl/2-propanol. The precipitate was filtered off and dried (vacuum; 80 °C), yielding 1.40 g (35%) of 4-amino-5-chloro-2,3-dihydro-N-[1-(3-methoxypropyl)-4-piperidinyl]-7-benzofurancarboxamide monohydrochloride (comp. 1).

Example 3

- Reaction under N₂ flow. 4-Amino-5-chloro-2,3-dihydro-7-benzofurancarboxylic acid (0.18 mol) was dissolved in tetrahydrofuran (360 ml) and this solution was stirred and cooled to ± 3 °C. 1,1'-Carbonylbis-1H-imidazole (0.18 mol) was added in one portion and cooling was stopped. The mixture was stirred for 75 minutes (became homogeneous after 30 minutes). A solution of 1-(3-methoxypropyl)-4-piperidinamine (0.18 mol) in tetrahydrofuran (90 ml) was added dropwise (exothermic temperature rise from 23 °C to 27 °C). The reaction mixture was stirred for 24 hours. More 1,1'-Carbonylbis-1H-imidazole (0.0125 mol) was added and the reaction mixture was stirred for 75 minutes. More 1-(3-methoxypropyl)-4-piperidinamine (0.0125 mol) was added (in 10 ml THF). The resulting reaction mixture was stirred for 3 hours at room temperature, then for 2.5 hours at reflux temperature. Then, the mixture was stirred for 13 hours, allowing it to cool to room temperature. The solvent was evaporated. The residue was stirred for 8 hours in water (360 ml) and the precipitate was filtered off, washed with water, then dried (vacuum; 30 °C), yielding 62.9 g (95%) 4-amino-5-chloro-2,3-dihydro-N-[1-(3-methoxypropyl)-4-piperidinyl]-7-benzofurancarboxamide monohydrate; mp. 90.7 °C (comp. 2).

Example 4

- Compound (2) (5 g, 0.0129 mol) was dissolved in warm ethanol (25 ml). A solution of (+)-(S)-lactic acid (1.45 g, 0.0135 mol) in ethanol (10 ml) was added. Under continuous stirring, crystallization started at 23 °C. The mixture was stirred for 24 hours. The precipitate was filtered off, washed with ethanol (2 ml), then dried (vacuum; 55 °C; 72 hours), yielding 3.7 g (62%) of 4-amino-5-chloro-2,3-dihydro-N-[1-(3-methoxypropyl)-4-piperidiny]-7-benzofurancarboxamide (+)-(S)-2-hydroxypropanoic acid salt (1:1); mp. 170.4°C (comp. 3).
- Compound (2) (5 g, 0.0129 mol) was dissolved in warm ethanol (35ml)/water (3.5 ml). Phosphoric acid (0.929 ml) was added and crystallization almost immediately resulted. The mixture was stirred for 24 hours at 23 °C. The precipitate was filtered off, washed with ethanol (2 ml), then dried (vacuum; 55 °C; 72 hours), yielding 5.87 g (97.7%) 4-amino-5-chloro-2,3-dihydro-N-[1-(3-methoxypropyl)-4-piperidiny]-7-benzofurancarboxamide phosphoric acid salt (1:1); mp. 259.6°C (comp. 4).
- Compound (2) (5 g, 0.0129 mol) was dissolved in warm ethanol (35 ml)/water (3.5 ml). A 48% hydrobromic acid solution (1.52 ml) was added and crystallization almost immediately resulted. The mixture was stirred for 24 hours at 13 °C. The precipitate was filtered off, washed with ethanol (2 ml), then dried (vacuum; 55 °C; 72 hours), yielding 5.4 g (93.2%) of 4-amino-5-chloro-2,3-dihydro-N-[1-(3-methoxypropyl)-4-piperidiny]-7-benzofurancarboxamide . hydrobromide (1:1); mp. 280.1°C (comp. 5).
- Compound (2) (5 g, 0.0129 mol) was dissolved in warm ethanol (25 ml). A solution of succinic acid (1.6 g) in ethanol (10 ml)/water (3.5 ml) was added. Upon scratching, crystallization resulted. The mixture was stirred for 24 hours at 23 °C. The precipitate was filtered off, washed with ethanol (2 ml), then dried (vacuum; 55 °C; 72 hours), yielding 5.7 g (91%) of 4-amino-5-chloro-2,3-dihydro-N-[1-(3-methoxypropyl)-4-piperidiny]-7-benzofurancarboxamide . butanedioate (1:1); mp. 197.2°C (comp. 6).

Example 5

- Compound (2) (5 g, 0.0129 mol) was dissolved in ethanol (35 ml). Water (3.5 ml) was added. Sulfuric acid (0.75 ml) was added dropwise. The mixture was stirred for 24 hours at ± 22°C. The precipitate was filtered off, washed with ethanol (2 ml), then dried (vacuum; 55-60 °C; 72 hours), yielding 6.1 g (101%) of 4-amino-5-chloro-2,3-dihydro-N-[1-(3-methoxypropyl)-4-piperidiny]-7-benzofurancarboxamide . sulfate (1:1); mp. 267.5°C (comp. 7).
- Compound (2) (5 g, 0.0129 mol) was dissolved in ethanol (35 ml). Water (3.5 ml) was added. Methanesulfonic acid (0.88 ml) was added dropwise. The mixture was stirred then dried (vacuum; 55-60 °C; 72 hours), yielding 6 g (100%) 4-amino-5-chloro-2,3-

dihydro-N-[1-(3-methoxypropyl)-4-piperidinyl]-7-benzofurancarboxamide .
methanesulfonic acid salt (1:1); mp. 286°C (comp. 8).

Compound (2) (5 g, 0.0129 mol) was dissolved in methylisobutyl ketone (35 ml), at 60-65 °C. Acetic acid (0.8 ml) was added dropwise (temperature rise to 75 °C).

- 5 Precipitation almost immediately resulted. The mixture was allowed to cool to room temperature. The mixture was stirred for 20 hours. The precipitate was filtered off, washed with ethanol (2 ml), then dried (vacuum; 55-60 °C; 72 hours), yielding 5.5 g (99%) of 4-amino-5-chloro-2,3-dihydro-N-[1-(3-methoxypropyl)-4-piperidinyl]-7-benzofurancarboxamide . acetic acid salt (1:1); mp. 156.1°C (comp. 9).

10

Pharmacological examples

Example 6

- Stimulation of non-adrenergic non-cholinergic nerves elicits relaxation followed by a contraction. The relaxation is mediated via a transmitter different from noradrenaline,
15 nitric oxide or ATP. The contraction is mediated via a transmitter different from acetylcholine.

- Dunkin-Hartley guinea pigs of either sex (350-600 g, not fasted) were killed by cervical dislocation followed by decapitation. The colon ascendens was removed and the lumen
20 was cleansed by repeated washing with De Jalon solution. After carefully dissecting the mesentery, the colon ascendens was divided into 4 segments of 3 cm length. Each segment was mounted vertically in an organ bath containing 100 ml De Jalon solution. The organ bath was kept at 37 °C and gassed with a mixture of 95 % oxygen and 5 % carbondioxide. In order to block the α , β , and muscarinic receptors, phentolamine (10^{-6}
25 M), propranolol (3×10^{-7} M) and atropine (3×10^{-7} M) were added to the solution. Contractions were measured isometrically. The preparation was repeatedly stretched until a basal tension of 40 mN was obtained and allowed to stabilize for 45 to 60 minutes. Histamine (3×10^{-5} M) was added to the bath solution in order to obtain a maximal contraction. Transmural excitation was applied over the whole length of the
30 colon strip by means of two platinum electrodes, the anode threaded through the lumen of the colon; the cathode in the bathing solution. The preparation was excited with rectangular square wave pulses (9 V, 1 ms/pulse) for 10 seconds every 5 minutes at different frequencies. Electrical stimulation resulted in a relaxation (= ON response) immediately followed by a contraction (= OFF response). Initially, the preparations
35 were stimulated three times at 0.4 Hz in order to obtain a submaximal relaxation, followed by three stimuli at 1.5 Hz in order to obtain a submaximal contraction. Then the test compound was added to the bath fluid and again both stimuli (0.4 Hz and 1.5 Hz)

were repeated three times.

At a concentration of 3×10^{-7} M the test compound induced an increase of the OFF response of 100% of initial value.

5 Example 7

Dunkin-Hartley guinea pigs of either sex (350 g or more, not fasted) were killed by cervical dislocation followed by decapitation. The colon descendens was cut at ± 5 cm from the rectum, cut and ligated at a length of ± 40 cm and freed of adhering tissue.

10 When there were at least 10 pellets in the colon, the tissue was transferred to a glass beaker containing 200 ml of Krebs-Henseleit solution, gassed with a mixture of 95 % oxygen and 5 % carbondioxide and maintained at 37 °C. The solution contained either pure solvent or the test compound. The expelled pellets were counted and removed from the solution every 5 minutes during a maximum period of 60 minutes.

15 The cumulative number of pellets expelled from the colon at every point was expressed as the percentage of the total number of pellets present in the entire colon at the start of the experiment. Time response curves were made by plotting the cumulative percentage of pellets expelled from the colon versus time.

20 At a concentration of 3×10^{-9} M of the present compound 80 % of the initial amount of pellets was expelled within 10 minutes.

Example 8


Guinea Pig Ileum Coaxial Stimulation.

25 Dunkin Hartley guinea-pigs of both sexes (body weight ± 500 g) were killed by cervical dislocation followed by decapitation. The ileum was removed and cleansed with warmed and oxygenated Krebs-Henseleit solution. Non-terminal, intact ileum segments, 4.5 cm long, of the guinea pig were vertically suspended with a preload of 1 g in 100 ml Krebs-Henseleit solution (37.5°C), gassed with a mixture of 95% O₂ and 5% CO₂. Transmural excitation was applied over the whole length of the ileum segment by means of two
30 platinum electrodes, the anode threaded through the lumen of the ileum, the cathode in the bathing solution. The preparation was excited with single rectangular stimuli [1 msec; 0.1 Hz; submaximal response (current leading to 80% of maximal reponse)] from a programmable stimulator. Contractions were measured isometrically. During the stabilization period of 30 min, the strips were repeatedly stretched to a tension of 2g, in
35 order to obtain a steady state tension of 1g. Before starting the electrical stimulation, a cumulative dose response curve of acetylcholine was given. The electrical stimulation was started at supramaximal current to determine the maximal amplitude of the twitch

- responses. When these responses were stable, a submaximal stimulation to obtain 80% of the maximal responses was given until the twitch responses were constant for at least 15 min, whereafter a single dose of the test compound was added to the bath fluid. The amplitude of the twitch response five minutes after the administration of the test
- 5 compound is compared with the amplitude before the administration of the test compound. The present compound showed an increase of the amplitude of the twitch response of more than 5% at a concentration of $3 \cdot 10^{-9}$ M.

Claims

1. A compound of formula



or a pharmaceutically acceptable acid addition salt thereof.

- 10
2. A compound as claimed in claim 1 wherein the compound is 4-amino-5-chloro-2,3-dihydro-N-[1-(3-methoxypropyl)-4-piperidinyl]-7-benzofurancarboxamide monohydrochloride.
- 15
3. A compound as claimed in claim 1 wherein the compound is 4-amino-5-chloro-2,3-dihydro-N-[1-(3-methoxypropyl)-4-piperidinyl]-7-benzofurancarboxamide butanedioate (1:1).
- 20
4. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound as claimed in any of claims 1 to 3.
- 25
5. A process of preparing a pharmaceutical composition as claimed in claim 4 wherein a therapeutically effective amount of a compound as claimed in any of claims 1 to 3 is intimately mixed with a pharmaceutically acceptable carrier.
- 30
6. A compound as claimed in any of claims 1 to 3 for use as a medicine.
7. The use of a compound as claimed in any of claims 1 to 3 for the manufacture of a medicament for treating gastrointestinal motility disorders.
8. The use of a compound as claimed in any of claims 1 to 3 for the manufacture of a medicament for treating constipation.
9. The use of a compound as claimed in any of claims 1 to 3 for the manufacture of a medicament for treating irritable bowel disease.
10. The use of a compound as claimed in any of claims 1 to 3 for the manufacture of a

-12-


medicament for treating intestinal atony.

11. The use of a compound as claimed in any of claims 1 to 3 for the manufacture of a medicament for accelerating (large) bowel transit.


5

12. A process of preparing a compound as claimed in claims 1 or 2 wherein
a) an intermediate of formula (II) is N-alkylated with an alkylating reagent of formula (III), wherein W is an appropriate leaving group such as a halo, e.g. chloro, or a sulfonyloxy leaving group in a reaction inert solvent in the presence of a suitable base;

10



- b) an amine of formula (V) is N-acylated with a carboxylic acid of formula (IV)



15

and, if desired, converting a compound of formula (I) into a therapeutically active non-toxic acid addition salt, or conversely, converting an acid addition salt into a free base form with alkali.

20

INTERNATIONAL SEARCH REPORT

Intern Application No

PCT/EP 95/04519

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 6 C07D405/12 K31/445

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 389 037 (JANSSEN PHARMACEUTICA N.V.) 26 September 1990 cited in the application see pages 27, 31, example 59 and claims. ---	1-12
Y	EP,A,0 445 862 (JANSSEN PHARMACEUTICA N.V.) 11 September 1991 cited in the application see pages 33, 34, example 107 and claims. -----	1-12

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

5 February 1996

Date of mailing of the international search report

14.02.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+ 31-70) 340-3016

Authorized officer

Chouly, J

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 95/04519

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-389037	26-09-90	AT-T- 128132	15-10-95
		AU-B- 616838	07-11-91
		AU-B- 5209190	27-09-90
		CA-A- 2012432	22-09-90
		CN-A- 1045781	03-10-90
		DE-D- 69022453	26-10-95
		FI-A- 944076	05-09-94
		IL-A- 93817	30-03-95
		IL-A- 110397	26-05-95
		JP-A- 2289566	29-11-90
		NO-B- 176101	24-10-94
		US-A- 5374637	20-12-94
EP-A-445862	11-09-91	AU-B- 636012	08-04-93
		AU-B- 7207991	12-09-91
		BG-B- 60381	31-01-95
		CA-A- 2037575	07-09-91
		CN-A- 1054598	18-09-91
		CN-A- 1054778	25-09-91
		JP-A- 4211685	03-08-92
		LT-A- 846	27-02-95
		LV-B- 10085	20-02-95
		NO-B- 177424	06-06-95
		US-A- 5185335	09-02-93
		US-A- 5262418	16-11-93